

BRAUERIA (Lunz am See, Austria) 36:4-10 (2009)

## Illustrations for several previously un-associated Arizona Trichoptera females.

David E. RUITER & Dean W. BLINN

**Abstract.** Genitalic illustrations are provided for 24 Arizona Trichoptera females. Male figures are also provided for *Clistoronia formosa*, *Clistoronia maculata* and *Psychoglypha schuhi*.

### Key Words:

Arizona, Trichoptera, females, illustrations.

Recent investigations of Arizona Trichoptera resulted in a preliminary list of 135 species reported from the state (BLINN & RUITER 2005). These collections included many insufficiently illustrated or un-associated females. Illustrations for 24 species are provided to improve our ability to conduct faunal surveys and immature associations.

### Materials and Methods

Adult collections were made primarily by light trap augmented by sweep net, examining spider webs and bridge supports, and larval rearing. All material used for illustrations was cleared with KOH. This paper is arranged alphabetically by family, genus and species. Three families (Apataniidae, Calamoceratidae, Leptoceridae) are briefly treated to provide the reader with updated references for available figures. The Glossosomatidae, Helicopsychidae, Hydroptilidae, Uenoidae, and Xiphocentronidae will be treated in separate publications.

#### Apataniidae

WIGGINS (1973) provides figures for the single apataniid (*Apatania arizona*) known from Arizona.

#### Brachycentridae

Two brachycentrid genera, with 5 species, are currently reported from Arizona (BLINN & RUITER 2005). Illustrations of the *Brachycentrus* females and the female of *Micrasema bacro* are provided by SCHMID (1983). ROSS (1947) indicated the close relationship between *M. onisca* and *M. arizonica*. CHAPIN (1978) indicated the *M. arizonica* type genitalia are missing but also indicated *M. arizonica* was probably distinct from *M. onisca*. We did not collect specimens which could be attributed to *M. arizonica*. The remains of the *M. arizonica* type were compared with a *M. onisca* paratype and distinguishable characters could not be found. The *M. onisca* female figure (fig. 1) is based on several male/female collections which were also compared with a *M. onisca* paratype. The female of *M. onisca* can be distinguished from *M. bacro* by the mesally divided 8<sup>th</sup> sternite of *M. onisca*. Arizona brachycentrid collections are from small, cool streams at altitudes >1550 m with low mean channel embeddedness (9.6%,  $\pm 2.0$ ; BLINN & RUITER 2005, 2006).

#### Calamoceratidae

While BLINN & RUITER (2005) was in press, the Arizona *Phylloicus* were treated as a single species (*P. aeneus*). PRATHER (2003) revised *Phylloicus* and clarified the distinction between *P. aeneus* and *P. mexicanus* resulting in two *Phylloicus* species in Arizona. PRATHER (2003) also provided female illustrations. Arizona *Phylloicus* collections are from small cool streams at altitudes between 1200-1750 m with a mean channel embeddedness of 20%,  $\pm 7.8$  (BLINN & RUITER 2005, 2006).

#### Hydrobiosidae

SCHMID (1989) indicated the close similarity between *Atopsyche* females and that conclusion is supported by the two Arizona species. Female *A. sperryi* (fig. 2) and *A. tripunctata* (fig. 3) can be separated by the shape of the lateral invagination of the 8<sup>th</sup> tergite along with other minor genitalic variations. The invagination of *A. sperryi* is globular in dorsal view, while that of *A. tripunctata* is more elongate and slightly bi-lobed. The species are sympatric with overlapping emergence periods, particularly in the Oak Creek drainage. When sympatric, *A. sperryi* is collected much more frequently. Arizona *Atopsyche* collections are from altitudes between 1346-2757 m. *Atopsyche sperryi* were found in streams with a lower mean channel embeddedness (1.9%,  $\pm 3.2$ ) than *A. tripunctata* (19.3%,  $\pm 5.7$ ) (BLINN & RUITER 2005, 2006).

#### Hydropsychidae

Fourteen species in four hydropsychid genera have been reported from Arizona (BLINN & RUITER 2005). SCHEFTER (2005) concluded that *Ceratopsyche* should be left at the subgenus level, reducing the Arizona hydropsychid genera to three. GORDON (1974) provides illustrations for the Arizona *Cheumatopsyche* females. FLINT (1974) provides illustrations for the Arizona *Smicridea* females. ROSS (1944) provides a female illustration for *Hydropsyche auricolor* (as *H. solex*). NIMMO (1987) contains a female illustration of *H. occidentalis*. Since preparation of BLINN & RUITER (2005) we have collected *Hydropsyche vanaca* (fig. 4) in Arizona. *Hydropsyche osleri* is readily distinguished from the other Arizona *Hydropsyche* species by the long clasper receptacle in lateral view (see NIMMO 1987). *H. vanaca* and *H. venada* (fig. 6) females are quite distinct but, in Arizona, possibly confused with *H. cockerelli* (see SCHEFTER et. al. 1986). In lateral and dorsal view the clasper receptacle of *H. vanaca* females is much larger than that of *H. venada* & *H. cockerelli*. The clasper receptacle of *H. venada*, while small in lateral view, appears as a long sinuate sclerotized area in dorsal view. The clasper receptacle of *H. cockerelli* is much shorter in dorsal view. Arizona hydropsychid collections occurred in a variety of streams. Typically, *H. venada* were found in streams at high altitudes with low channel embeddedness (10.3%,  $\pm 2.6$ ), while *Cheumatopsyche enonis*, *Smicridea dispar*, *S. fasciatella*, and *S. signata* were associated with streams with high mean channel embeddedness (53%,  $\pm 9.4$ ) over a wide range of altitudes (BLINN & RUITER 2005, 2006).

#### Lepidostomatidae

Female figures for six of the nine *Lepidostoma* species reported from Arizona are provided by WEAVER (1988). DENNING (1949) provided a brief description of the *L.*

*aporum* female and a figure of the vaginal apparatus. WEAVER (1988) did not have *L. aporum* females and noted the similarity between DENNING's figure and *L. rayneri*. *Lepidostoma aporum* females (fig. 6) can be readily distinguished from the other known *pluviale* group females by the upward directed ventromesal band in ventral view. There are two *Lepidostoma* (*L. apache* & *L. lacinatum*) for which we have yet to develop positive associations. We are providing illustrations of the vaginal apparatus for two un-associated Arizona *Lepidostoma* species (figs. 7 & 8) which may represent the unknown females. Arizona collections occurred in streams at 1,545 m with a low mean channel embeddedness 14.5%,  $\pm$  4.4 (BLINN & RUITER 2005, 2006).

#### Leptoceridae

With the exception of the five *Nectopsyche* females, female figures for the other three Arizona leptocerid genera, with seven species, are available (*O. arizonica* in DENNING 1951; *O. avara* and *O. disjuncta* in SMITH & LEHMKUHL 1980; *O. inconspicua* in ROSS 1944; *O. metlacensis* in BUENO-SORIA 1981; *Triaenodes tardus* and *Ylodes reuteri* in ROSS 1944). Characters to separate the Arizona *Nectopsyche* females were not discovered. Arizona leptocerid collections occurred over a wide range of altitudes in streams with a mean channel embeddedness of 24%  $\pm$  6.2. (BLINN & RUITER 2005, 2006).

#### Limnephilidae

Limnephilidae is the second most diverse family, behind the Hydroptilidae, in Arizona. Females figures are available for the Arizona species of *Amphicosmoecus* and *Anabolia* (see NIMMO 1971); *Hesperophylax* (see PARKER & WIGGINS 1985); *Limnephilus* (see RUITER 1995, RUITER 2007); *Onocosmoecus* (see WIGGINS & RICHARDSON 1986); and *Psychoglypha subborealis* (see SCHMID 1952). Arizona limnephilid collections occurred at elevations >1600 m with a mean channel embeddedness of 10%  $\pm$  1.8 (BLINN & RUITER 2005, 2006).

ESSIG (1926) reported *Clistoronia formosa* from Arizona without locality information. We have not collected additional specimens in Arizona. *Clistoronia maculata* is relatively common in Arizona at higher elevations. Recent figures for neither *C. formosa* or *C. maculata* are available so we have included male and female figures for both species (figs. 9 and 10).

Records for *Psychoglypha schuhi* seem to include only the Nevada types and a single Wyoming collection (RUITER 1999) available. We were surprised to find this species present near Greer. We have included both male and female figures (fig. 11).

#### Odontoceridae

*Marilia nobsca* was described from a female, although a figure was not provided. The two North American species are readily separated by spur count (*M. flexuosa*: 2-4-2; *M. nobsca*: 2-4-4). *Marilia flexuosa* is also about half the size of *M. nobsca*. SCHMID (1980) provided a *M. flexuosa* female figure. We have included a figure of the *M. nobsca* vaginal apparatus (fig. 12) for comparison. Arizona collections occurred at altitudes between 1085-2525 m with low mean channel embeddedness (5%,  $\pm$  0.1) (BLINN & RUITER 2005, 2006).

#### Philopotamidae

Arizona philopotamid collections occurred over a wide range of altitudes (640-2490 m) with a mean channel embeddedness of 17% ( $\pm$  4.5) (BLINN & RUITER 2005, 2006).

*Chimarra* are common in Arizona with seven species (*C. adella*, *C. angustipennis*, *C. primula*, *C. ridleyi*, *C. schiza*, *C. texana*, *C. utahensis*) occurring primarily in small streams and springs. Often two or three species are sympatric, and emerge at the same time. The Arizona females can be readily separated by the ventral view of the 8<sup>th</sup> & 9<sup>th</sup> segments (figs. 13 thru 19) *Chimarra texana* is unusual with an anteroventral spur on the 8<sup>th</sup> sternite. *Chimarra angustipennis* and *C. primula* are most likely confused and can be separated by the relative length of the 9<sup>th</sup> sternites.

To date only *Wormaldia arizonensis* has been reported from Arizona. The female is readily distinguished by the inverted v-shaped medial fold of the 8<sup>th</sup> sternite (fig. 20). The medial fold is covered with fine, dark setae.

#### Polycentropodidae

The two Arizona polycentropodid genera (*Polycentropus arizonensis*, *P. aztecus*, *P. gertschi*, *P. halidus*, *P. variegatus*, *Polypsectropus charlesi*) occur at elevations between 575-2600 m with a mean channel embeddedness of 14.2%  $\pm$  3.7 (BLINN & RUITER 2005, 2006).

Females of *P. halidus* and *P. variegatus* were figured by NIMMO (1986). Figures of *P. arizonensis* and *P. gertschi* are provided here (figs. 21 and 22). The *P. arizonensis* female can be readily separated from the known Arizona females by the narrow 8<sup>th</sup> sternal lateral lobes. It may be confused with *P. gertschi*, which also has relatively narrow 8<sup>th</sup> sternal lateral lobes, but the apex of the *P. arizonensis* vaginal apparatus has four distinct subparallel apical lobes while the apex of the *P. gertschi* vaginal apparatus is triangular in shape without distinct lobes.

A figure of the *Polypsectropus charlesi* female is also provided (fig. 23).

#### Sericostomatidae

Two species of *Gumaga* are reported from Arizona (*Gumaga griseola* & *Gumaga nigricula*). WOOD & RESH (1991) documented the variability within the two species and their inability to clearly distinguish the females. We have yet to collect Arizona males attributable to *G. nigricula*. A figure of the *G. griseola* female vaginal apparatus is provided (fig. 24) to possibly assist in separating these species. Collections of *G. griseola* occurred at elevations between 1618-2797 m with a mean channel embeddedness of 7.6%  $\pm$  2.6 (BLINN & RUITER 2005, 2006).

#### Acknowledgement:

We thank Oliver S. Flint, Jr. for assistance with several male/female associations.

#### Literature Cited:

BLINN, D.W. & D.E. RUITER. 2005. Caddisfly (Trichoptera) community structure and distribution in Arizona, USA: Effects of selected environmental determinants in Proceedings of the 11<sup>th</sup> International Symposium on

- Trichoptera (2003, Osaka). Tokai University Press, Osaka. Japan. Pages 63-71.
- BLINN, D.W. & D.E. RUITER. 2006. Tolerance values of stream caddisflies (Trichoptera) in the lower Colorado River Basin, USA. *Southwestern Naturalist* 51(3):326-337.
- BUENO-SORIA, J. 1981. Estudios en insectos acuáticos de México I. Trichoptera (Leptoceridae). Cinco nuevas especies de *Oecetis* MCLACHLAN. *Folia Entomológica Mexicana*, 49:103-120.
- CHAPIN, J. W. 1978. Systematics of Nearctic *Micrasema* (Trichoptera: Brachycentridae). PhD Thesis, Clemson University, 136 pages.
- DENNING, D.G. 1949. New and little known species of caddis flies. *American Midland Naturalist* 42(1): 112-122.
- ESSIG, E.O. 1926. Insects of Western North America, The MacMillan Company, New York, 1035 pages.
- FLINT, O.S. 1974. Studies of Neotropical caddisflies, XVII: The genus *Smicridea* from north and central America (Trichoptera: Hydropsychidae). *Smithsonian Contributions to Zoology* 175. 65 pages.
- GORDON, A.E. 1974. A synopsis and phylogenetic outline of the Nearctic members of Cheumatopsyche. *Proceedings of the Academy of Natural Sciences Philadelphia* 126(9):117-160.
- NIMMO, A.P. 1971. The adult Rhyacophilidae and Limnephilidae (Trichoptera) of Alberta and eastern British Columbia, and their post-glacial origins. *Quaestiones Entomologicae* 7:3-234.
- NIMMO, A.P. 1986. The adult Polycentropodidae of Canada and adjacent United States. *Quaestiones Entomologicae*. 22: 143-252.
- NIMMO, A.P. 1987. The adult Arctopsychoidea and Hydropsychidae (Trichoptera) of Canada and adjacent United States. *Quaestiones Entomologicae* 23:1-189.
- PARKER, C.R., and G.B. WIGGINS. 1985. The Nearctic caddisfly genus *Hesperophylax* (Trichoptera: Limnephilidae). *Canadian Journal of Zoology* 61: 2443-2472.
- PRATHER, A.L. 2003. Revision of the Neotropical caddisfly genus *Phylloicus* (Trichoptera: Calamoceratidae). *Zootaxa* 275:1-214.
- ROSS, H.H. 1944. The caddis flies or Trichoptera of Illinois. *Bulletin of the Illinois Natural History Survey* 23(1):1-326.
- ROSS, H.H. 1947. Descriptions and records of North American Trichoptera with synoptic notes. *Transactions of the American Entomological Society* 73:125-168.
- RUITER, D.E. 1999. A new species and a new synonym in the genus *Psychoronia* (Limnephilidae), with significant records for caddisflies from western North America (Trichoptera). *Great Basin Naturalist* 59(2):160-168.
- RUITER, D.E. 2007. Description of the *Limnephilus granti* (Nimmo) female with a redescription of the male (Trichoptera: Limnephilidae). *Proceedings of the Entomological Society of Washington* 109(1):86-89.
- SCHEFTER, P.W. 2005. Re-evaluation of genera in the subfamily Hydropsychinae (Trichoptera: Hydropsychidae). *Aquatic Insects* 27(2):133-154.
- SCHEFTER, P.W., G.B. WIGGINS and J.D. UNZICKER. 1986. A proposal for assignment of *Ceratopsyche* as a subgenus of *Hydropsyche*, with new synonyms and a new species (Trichoptera: Hydropsychidae). *Journal of the North American Benthological Society* 5(1):67-84.
- SCHMID, F. 1952. Le groupe de *Chilostigma* (Trichopt., Limnoph.). *Archive für Hydrobiologie* 47(1):75-163.
- SCHMID, F. 1980. Genera des Trichoptères du Canada et des États adjacents. *Les Insectes et Arachnides du Canada. Partie 7. Agriculture Canada Publications* 1692.
- SCHMID, F. 1983. Revision des Trichoptères Canadiens . III. Les Hyalopsychidae, Psychomyiidae, Goeridae, Brachycentridae, Sericostomatidae, Helicopsychidae, Beraeidae, Odontoceridae, Calamoceratidae et Molannidae. *Mém. de la Société Entomologique du Canada*, No. 125.
- SCHMID, F. 1989. Les Hydrobiosides (Trichoptera, Annulipalpia). *Bulletin de l'Institut Royal des Sciences Naturelles de Belgique, Entomologie Vol.59 Supplement*, 154 pages.
- SMITH, D.H. and D.M. LEHMKUHL. 1980. Analysis of two problematic North American caddisfly species: *Oecetis avara* (Banks) and *Oecetis disjuncta* (Banks) (Trichoptera: Leptoceridae) *Quaestiones Entomologicae* 16:641-656.
- WEAVER, J.S. 1988. A synopsis of the North American Lepidostomatidae (Trichoptera). *Contributions of the American Entomological Institute* 24(2):1-141.
- WIGGINS, G.B. 1973. Contributions to the systematics of the caddisfly family Limnephilidae (Trichoptera). I. *Royal Ontario Museum, Life Sciences Contributions* 94.
- WIGGINS, G.B. and J.S. RICHARDSON. 1986. Revision of the *Onocosmoecus unicolor* group (Trichoptera: Limnephilidae, Dicosmoecinae). *Psyche* 93(3-4):187-216.
- WOOD, J.R. and V.H. RESH. 1991. Morphological and ecological variation in stream and spring populations of *Gumaga nigracula* (MCLACHLAN) in the California (USA) coast ranges. *Proceedings of the Sixth International Symposium on Trichoptera. Adam Mickiewicz University Press. Pages* 14-20.
- Authors:** David E. Ruiter, 6260 S. Grant Street, Centennial, CO, USA 80121. - Dr. Dean W. Blinn, Northern Arizona University, Flagstaff, AZ, USA 86011-5640




Fig. 1. *Micrasema onisca* female: (a) apical abdominal segments, lateral view; (b) apical abdominal segments, ventral view; (c) vaginal apparatus, ventral view. - Fig. 2. *Atopsyche sperryi* female: (a) apical abdominal segments, dorsal view; (b) vaginal apparatus, ventral view. - Fig. 3. *Atopsyche tripunctata* female: (a) apical abdominal segments, lateral view; (b) vaginal apparatus, ventral view. - Fig. 4. *Hydropsyche vanaca* female: (a) apical abdominal segments, lateral view; (b) apical abdominal segments, dorsal view. - Fig. 5. *Hydropsyche venada* female: (a) apical abdominal segments, lateral view; (b) apical abdominal segments, dorsal view. - Fig. 6. *Lepidostoma apornum* female: (a) apical abdominal segments, lateral view; (b) vaginal apparatus, ventral view. - Fig. 7. *Lepidostoma* species 1 female: (a) vaginal apparatus, lateral view; (b) vaginal apparatus, ventral view. - Fig. 8. *Lepidostoma* species 2 female: (a) vaginal apparatus, lateral view; (b) vaginal apparatus, ventral view.




Fig. 9. *Clistoronia formosa* adults: (a) male apical abdominal segments, lateral view; (b) male apical abdominal segments, dorsal view; (c) male apical abdominal segments, caudal view; (d) male aedeagus, lateral view; (e) female apical abdominal segments, lateral view; (f) female apical abdominal segments, dorsal view; (g) female apical abdominal segments, ventral view. - Fig. 10. *Clistoronia maculata* adults: (a) male apical abdominal segments, lateral view; (b) male apical abdominal segments, dorsal view; (c) male apical abdominal segments, caudal view; (d) male aedeagus, lateral view; (e) female apical abdominal segments, lateral view; (f) female apical abdominal segments, dorsal view; (g) female apical abdominal segments, ventral view.




Fig. 11. *Psychoglypha schuhi* adults: (a) male apical abdominal segments, lateral view; (b) male apical abdominal segments, dorsal view; (c) male apical abdominal segments, ventral view; (d) male aedeagus, lateral view; (e) male aedeagus, dorsal view; (f) female apical abdominal segments, lateral view; (g) female apical abdominal segments, dorsal view; (h) female apical abdominal segments, ventral view; (i) female vaginal apparatus, ventral view. - Fig. 12. *Marilia nobisca* female: female vaginal apparatus, ventral view. - Fig. 13. *Chimarra adella* female: apical abdominal segments, ventral view. - Fig. 14. *Chimarra angustipennis* female: apical abdominal segments, ventral view. - Fig. 15. *Chimarra primula* female: apical abdominal segments, ventral view. - Fig. 16. *Chimarra ridleyi* female: apical abdominal segments, ventral view. - Fig. 17. *Chimarra schiza* female: apical abdominal segments, ventral view. - Fig. 18. *Chimarra texana* female: apical abdominal segments, ventral view. - Fig. 19. *Chimarra utahensis* female: apical abdominal segments, ventral view.




Fig. 20. *Wormaldia arizonensis* female: apical abdominal segments, ventral view. - Fig. 21. *Polycentropus arizonensis* female: apical abdominal segments, ventral view. - Fig. 22. *Polycentropus gertschi* female: apical abdominal segments, ventral view. - Fig. 23. *Polyplectropus charlesi* female: apical abdominal segments, ventral view. - Fig. 24. *Gumaga griseola* female: (a) vaginal apparatus, lateral view; (b) vaginal apparatus, ventral view.



### Change of addresses

Dénes SCHMERA  
Institute of Biology  
College of Nyíregyháza  
Sóstói út 31/B  
H – 4400 Nyíregyháza

Peter NEU  
Heiligenbungert 1  
D -54317 KASEL

Dr.Gisli M.GISLASON  
University of Iceland  
Askja- Natural Science Building  
Sturlugata 7  
IS - 101 REYKJAVIK

Dr.John B.WARD  
94c Simeon Street  
Spreydon  
CHRISTCHURCH 8024  
Neuseeland

Monika HESS  
Büro H2  
Corneliusstraße 30  
D - 80469 MÜNCHEN

Prof. Dr. János Oláh  
Tarján u. 28  
H – 4032 DEBRECEN  
profolah@gmail.com